arXiv:1303.2527v1 [math.CV] 11 Mar 2013

INITIAL COEFFICIENT BOUNDS FOR A GENERAL CLASS OF BI-UNIVALENT FUNCTIONS

H. ORHAN*, N. MAGESH, AND V.K.BALAJI

Abstract. Inspired by the recent works of Srivastava et al. [14], Frasin and Aouf [6] and others [1, 5, 7, 17, 18], we propose to investigate the coefficient estimates for a general class of analytic and bi-univalent functions. Also, we obtain estimates on the coefficients $|a_2|$ and $|a_3|$ for functions in this new class. Some interesting remarks, corollaries and applications of the results presented here are also discussed.

1. Introduction

Let $\mathcal{A}$ denote the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \tag{1.1}$$

which are analytic in the open unit disk $\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$. Further, by $\mathcal{S}$ we shall denote the class of all functions in $\mathcal{A}$ which are univalent in $\mathbb{U}$.

For two functions f and g, analytic in $\mathbb{U}$, we say that the function $f(z)$ is subordinate to $g(z)$ in $\mathbb{U}$, and write

$$f(z) \prec g(z), \quad z \in \mathbb{U},$$

if there exists a Schwarz function $w(z)$, analytic in $\mathbb{U}$, with

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1, \quad z \in \mathbb{U},$$

such that

$$f(z) = g(w(z)), \quad z \in \mathbb{U}.$$

In particular, if the function g is univalent in $\mathbb{U}$, the above subordination is equivalent to

$$f(0) = g(0) \quad \text{and} \quad f(\mathbb{U}) \subset g(\mathbb{U}).$$

Some of the important and well-investigated subclasses of the univalent function class $\mathcal{S}$ include (for example) the class $\mathcal{S}^*(\alpha)$ of starlike functions of order α in $\mathbb{U}$ and the class $\mathcal{K}(\alpha)$ of convex functions of order α in $\mathbb{U}$. By definition, we have

$$\mathcal{S}^*(\alpha) := \left\{ f : f \in \mathcal{S} \,\text{and}\, \Re\left(\frac{zf'(z)}{f(z)}\right) > \alpha;\ z \in \mathbb{U};\ 0 \le \alpha < 1 \right\} \tag{1.2}$$

and

$$\mathcal{K}(\alpha) := \left\{ f : f \in \mathcal{S} \,\text{and}\, \Re\left(1 + \frac{zf''(z)}{f'(z)}\right) > \alpha;\ z \in \mathbb{U};\ 0 \le \alpha < 1 \right\}. \tag{1.3}$$

2000 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Analytic functions, univalent functions, bi-univalent functions, starlike and convex functions, bi-starlike and bi-convex functions.

*Corresponding author e-mail: orhanhalit607@gmail.com.

It readily follows from the definitions (1.2) and (1.3) that

$$f \in \mathcal{K}(\alpha) \Longleftrightarrow zf' \in \mathcal{S}^*(\alpha).$$

Also, let $\mathcal{S}_{\mathcal{P}}^{\beta}(\alpha)$ and $\mathcal{C}_{\mathcal{P}}^{\beta}(\alpha)$ denote the subclasses of $\mathcal{S}$ consisting functions $f(z)$ which are defined, respectively by

$$\mathcal{S}_{\mathcal{P}}^{\beta}(\alpha) := \left\{ f : f \in \mathcal{S} \text{ and } \Re\left(e^{i\beta}\frac{zf'(z)}{f(z)}\right) > \alpha\cos\beta;\ z \in \mathbb{U};\ 0 \le \alpha < 1,\ \beta \in (-\frac{\pi}{2}, \frac{\pi}{2}) \right\} \tag{1.4}$$

and

$$\mathcal{C}_{\mathcal{P}}^{\beta}(\alpha) := \left\{ f : f \in \mathcal{S} \text{ and } \Re\left(e^{i\beta}\frac{z(f'(z))'}{f'(z)}\right) > \alpha\cos\beta;\ z \in \mathbb{U};\ 0 \le \alpha < 1,\ \beta \in (-\frac{\pi}{2}, \frac{\pi}{2}) \right\}. \tag{1.5}$$

It is easy to see that

$$f \in \mathcal{C}_{\mathcal{P}}^{\beta}(\alpha) \Longleftrightarrow zf' \in \mathcal{S}_{\mathcal{P}}^{\beta}(\alpha).$$

It is well known that every function $f \in \mathcal{S}$ has an inverse f^{-1}, defined by

$$f^{-1}(f(z)) = z,\ z \in \mathbb{U}$$

and

$$f(f^{-1}(w)) = w,\ |w| < r_0(f);\ r_0(f) \ge \frac{1}{4},$$

where

$$f^{-1}(w) = w - a_2w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2a_3 + a_4)w^4 + \dots. \tag{1.6}$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in $\mathbb{U}$ if both $f(z)$ and $f^{-1}(z)$ are univalent in $\mathbb{U}$. Let Σ denote the class of bi-univalent functions in $\mathbb{U}$ given by (1.1). Examples of functions in the class Σ are

$$\frac{z}{1-z},\ -\log(1-z),\ \frac{1}{2}\log\left(\frac{1+z}{1-z}\right)$$

and so on. However, the familiar Koebe function is not a member of Σ. Other common examples of functions in $\mathcal{S}$ such as

$$z - \frac{z^2}{2} \text{ and } \frac{z}{1-z^2}$$

are also not members of Σ (see [6, 14]).

In 1967, Lewin [9] investigated the bi-univalent function class Σ and showed that $|a_2| < 1.51$. On the other hand, Brannan and Clunie [2] (see also [3, 4, 15]) and Netanyahu [11] made an attempt to introduce various subclasses of the bi-univalent function class Σ and obtained non-sharp coefficient estimates on the first two coefficients $|a_2|$ and $|a_3|$ of (1.1). But the coefficient problem for each of the following Taylor-Maclaurin coefficients $|a_n|\,(n \in \mathbb{N} \setminus \{1,2\};\ \ \mathbb{N} := \{1,2,3,\cdots\})$ is still an open problem. Following Brannan and Taha [4], many researchers (see[1, 5, 6, 7, 8, 10, 13, 14, 17, 18]) have recently introduced and investigated several interesting subclasses of the bi-univalent function class Σ and they have found non-sharp estimates on the first two Taylor-Maclaurin coefficients $|a_2|$ and $|a_3|$.

Motivated by the above mentioned works, we define the following subclass of function class Σ.

Definition 1.1. Let $h : \mathbb{U} \to \mathbb{C}$, be a convex univalent function such that $h(0) = 1$ and $h(\bar{z}) = \overline{h(z)}$, for $z \in \mathbb{U}$ and $\Re(h(z)) > 0$. A function $f(z)$ given by (1.1) is said to be in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, h)$ if the following conditions are satisfied:

$$f \in \Sigma,\ e^{i\beta}\left((1-\lambda)\left(\frac{f(z)}{z}\right)^{\mu} + \lambda f'(z)\left(\frac{f(z)}{z}\right)^{\mu-1}\right) \prec h(z)\cos\beta + i\sin\beta, \quad z \in \mathbb{U} \quad (1.7)$$

and

$$e^{i\beta}\left((1-\lambda)\left(\frac{g(w)}{w}\right)^{\mu} + \lambda g'(w)\left(\frac{g(w)}{w}\right)^{\mu-1}\right) \prec h(w)\cos\beta + i\sin\beta, \quad w \in \mathbb{U}, \quad (1.8)$$

where $\beta \in (-\pi/2, \pi/2)$, $\lambda \geq 1$, $\mu \geq 0$ and the function g is given by

$$g(w) = w - a_2 w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2a_3 + a_4)w^4 + \dots \quad (1.9)$$

the extension of f^{-1} to $\mathbb{U}$.

Remark 1.2. If we set $h(z) = \frac{1+Az}{1+Bz}$, $-1 \leq B < A \leq 1$, in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, h)$, we have $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, \frac{1+Az}{1+Bz})$ and defined as

$$f \in \Sigma,\ e^{i\beta}\left((1-\lambda)\left(\frac{f(z)}{z}\right)^{\mu} + \lambda f'(z)\left(\frac{f(z)}{z}\right)^{\mu-1}\right) \prec \frac{1+Az}{1+Bz}\cos\beta + i\sin\beta, \quad z \in \mathbb{U}$$

and

$$e^{i\beta}\left((1-\lambda)\left(\frac{g(w)}{w}\right)^{\mu} + \lambda g'(w)\left(\frac{g(w)}{w}\right)^{\mu-1}\right) \prec \frac{1+Aw}{1+Bw}\cos\beta + i\sin\beta, \quad w \in \mathbb{U},$$

where $\beta \in (-\pi/2, \pi/2)$, $\lambda \geq 1$, $\mu \geq 0$ and the function g is given by (1.9).

Remark 1.3. Taking $h(z) = \frac{1+(1-2\alpha)z}{1-z}$, $0 \leq \alpha < 1$ in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, h)$, we have $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, \alpha)$ and $f \in \mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, \alpha)$ if the following conditions are satisfied:

$$f \in \Sigma,\ \Re\left(e^{i\beta}\left((1-\lambda)\left(\frac{f(z)}{z}\right)^{\mu} + \lambda f'(z)\left(\frac{f(z)}{z}\right)^{\mu-1}\right)\right) > \alpha\cos\beta,\ 0 \leq \alpha < 1;\ z \in \mathbb{U}$$

and

$$\Re\left(e^{i\beta}\left((1-\lambda)\left(\frac{g(w)}{w}\right)^{\mu} + \lambda g'(w)\left(\frac{g(w)}{w}\right)^{\mu-1}\right)\right) > \alpha\cos\beta,\ 0 \leq \alpha < 1;\ w \in \mathbb{U},$$

where $\beta \in (-\pi/2, \pi/2)$, $\lambda \geq 1$, $\mu \geq 0$ and the function g is given by (1.9).

Remark 1.4. Taking $\lambda = 1$ and $h(z) = \frac{1+(1-2\alpha)z}{1-z}$, $0 \leq \alpha < 1$ in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, h)$, we have $\mathcal{NP}_{\Sigma}^{\mu,1}(\beta, \alpha)$ and $f \in \mathcal{NP}_{\Sigma}^{\mu,1}(\beta, \alpha)$ if the following conditions are satisfied:

$$f \in \Sigma,\ \Re\left(e^{i\beta} f'(z)\left(\frac{f(z)}{z}\right)^{\mu-1}\right) > \alpha\cos\beta,\ 0 \leq \alpha < 1;\ \mu \geq 0;\ z \in \mathbb{U}$$

and

$$\Re\left(e^{i\beta} g'(w)\left(\frac{g(w)}{w}\right)^{\mu-1}\right) > \alpha\cos\beta,\ 0 \leq \alpha < 1;\ \mu \geq 0;\ w \in \mathbb{U},$$

where $\beta \in (-\pi/2, \pi/2)$ and the function g is given by (1.9).

Remark 1.5. Taking $\mu + 1 = \lambda = 1$ and $h(z) = \frac{1+(1-2\alpha)z}{1-z}$, $0 \le \alpha < 1$ in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, h)$, we have $\mathcal{NP}_{\Sigma}^{0,1}(\beta, \alpha)$ and $f \in \mathcal{NP}_{\Sigma}^{0,1}(\beta, \alpha)$ if the following conditions are satisfied:

$$f \in \Sigma,\ \Re\left(e^{i\beta}\frac{zf'(z)}{f(z)}\right) > \alpha\cos\beta,\ 0 \le \alpha < 1;\ z \in \mathbb{U}$$

and

$$\Re\left(e^{i\beta}\frac{wg'(w)}{g(w)}\right) > \alpha\cos\beta,\ 0 \le \alpha < 1;\ w \in \mathbb{U},$$

where $\beta \in (-\pi/2, \pi/2)$ and the function g is given by (1.9).

Remark 1.6. Taking $\mu = 1$ and $h(z) = \frac{1+(1-2\alpha)z}{1-z}$, $0 \le \alpha < 1$ in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, h)$, we have $\mathcal{NP}_{\Sigma}^{1,\lambda}(\beta, \alpha)$ and $f \in \mathcal{NP}_{\Sigma}^{1,\lambda}(\beta, \alpha)$ if the following conditions are satisfied:

$$f \in \Sigma,\ \Re\left(e^{i\beta}\left((1-\lambda)\frac{f(z)}{z} + \lambda f'(z)\right)\right) > \alpha\cos\beta,\ 0 \le \alpha < 1;\ \lambda \ge 1;\ z \in \mathbb{U}$$

and

$$\Re\left(e^{i\beta}\left((1-\lambda)\frac{g(w)}{w} + \lambda g'(w)\right)\right) > \alpha\cos\beta,\ 0 \le \alpha < 1;\ \lambda \ge 1;\ w \in \mathbb{U},$$

where $\beta \in (-\pi/2, \pi/2)$ and the function g is given by (1.9).

Remark 1.7. Taking $\mu = \lambda = 1$ and $h(z) = \frac{1+(1-2\alpha)z}{1-z}$, $0 \le \alpha < 1$ in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, h)$, we have $\mathcal{NP}_{\Sigma}^{1,1}(\beta, \alpha)$ and $f \in \mathcal{NP}_{\Sigma}^{1,1}(\beta, \alpha)$ if the following conditions are satisfied:

$$f \in \Sigma,\ \Re\left(e^{i\beta}f'(z)\right) > \alpha\cos\beta,\ 0 \le \alpha < 1;\ z \in \mathbb{U}$$

and

$$\Re\left(e^{i\beta}g'(w)\right) > \alpha\cos\beta,\ 0 \le \alpha < 1;\ w \in \mathbb{U},$$

where $\beta \in (-\pi/2, \pi/2)$ and the function g is given by (1.9).

We note that

1. $\mathcal{NP}_{\Sigma}^{1,1}(0, \alpha) = \mathcal{H}_{\Sigma}^{\alpha}$ (see [14])
2. $\mathcal{NP}_{\Sigma}^{1,\lambda}(0, \alpha) = \mathcal{B}_{\Sigma}(\alpha, \lambda)$ (see [6])
3. $\mathcal{NP}_{\Sigma}^{0,1}(0, \alpha) = \mathcal{F}_{\Sigma}(\alpha)$ (see [10])
4. $\mathcal{NP}_{\Sigma}^{\mu,1}(0, \alpha) = \mathcal{N}_{\Sigma}^{\mu}(\alpha)$ (see [13])
5. $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(0, \alpha) = \mathcal{N}_{\Sigma}^{\mu,\lambda}(\alpha)$ (see [5]).

In order to derive our main result, we have to recall here the following lemma.

Lemma 1.8. [12, 16] *Let the function* $\varphi(z)$ *given by*

$$\varphi(z) = \sum_{n=1}^{\infty} B_n z^n, \quad z \in \mathbb{U}$$

be convex in $\mathbb{U}$. *Suppose also that the function* $h(z)$ *given by*

$$h(z) = \sum_{n=1}^{\infty} h_n z^n, \quad z \in \mathbb{U}$$

is holomorphic in $\mathbb{U}$. *If* $h(z) \prec \varphi(z)$, $z \in \mathbb{U}$, *then* $|h_n| \le |B_1|$, $n \in \mathbb{N} = \{1, 2, 3, \dots\}$.

The object of the present paper is to introduce a general new subclass $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta,h)$ of the function class Σ and obtain estimates of the coefficients $|a_2|$ and $|a_3|$ for functions in this new class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta,h)$.

2. COEFFICIENT BOUNDS FOR THE FUNCTION CLASS $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta,h)$

In this section we find the estimates on the coefficients $|a_2|$ and $|a_3|$ for functions in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta,h)$.

Theorem 2.1. *Let $f(z)$ given by (1.1) be in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta,h)$, $0 \leq \alpha < 1$, $\lambda \geq 1$ and $\mu geq 0$, then*

$$|a_2| \leq \sqrt{\frac{2|B_1|\cos\beta}{(1+\mu)(2\lambda+\mu)}} \tag{2.1}$$

and

$$|a_3| \leq \frac{2|B_1| cos\beta}{(2\lambda+\mu)(1+\mu)}, \tag{2.2}$$

where $\beta \in (-\pi/2, \pi/2)$.

Proof. It follows from (1.7) and (1.8) that there exists $p, q \in \mathcal{P}$ such that

$$e^{i\beta}\left((1-\lambda)\left(\frac{f(z)}{z}\right)^{\mu} + \lambda f'(z)\left(\frac{f(z)}{z}\right)^{\mu-1}\right) = p(z)\cos\beta + i\sin\beta \tag{2.3}$$

and

$$e^{i\beta}\left((1-\lambda)\left(\frac{g(w)}{w}\right)^{\mu} + \lambda g'(w)\left(\frac{g(w)}{w}\right)^{\mu-1}\right) = p(w)\cos\beta + i\sin\beta, \tag{2.4}$$

where $p(z) \prec h(z)$ and $q(w) \prec h(w)$ have the forms

$$p(z) = 1 + p_1 z + p_2 z^2 + \dots, \ z \in \mathbb{U} \tag{2.5}$$

and

$$q(z) = 1 + q_1 w + q_2 w^2 + \dots, \ w \in \mathbb{U}. \tag{2.6}$$

Equating coefficients in (2.3) and (2.4), we get

$$e^{i\beta}(\lambda+\mu)a_2 = p_1\cos\beta \tag{2.7}$$

$$e^{i\beta}\left[\frac{a_2^2}{2}(\mu-1) + a_3\right](2\lambda+\mu) = p_2\cos\beta \tag{2.8}$$

$$-e^{i\beta}(\lambda+\mu)a_2 = q_1\cos\beta \tag{2.9}$$

and

$$e^{i\beta}\left[(\mu+3)\frac{a_2^2}{2} - a_3\right](2\lambda+\mu) = q_2\cos\beta. \tag{2.10}$$

From (2.7) and (2.9), we get

$$p_1 = -q_1 \tag{2.11}$$

and

$$2e^{i\beta}(\lambda+\mu)^2 a_2^2 = (p_1^2 + q_1^2)\cos^2\beta. \tag{2.12}$$

Also, from (2.8) and (2.10), we obtain

$$a_2^2 = \frac{e^{-i\beta}(p_2+q_2)\cos\beta}{(1+\mu)(2\lambda+\mu)}. \tag{2.13}$$

Since $p, q \in h(\mathbb{U})$, applying Lemma 1.8, we immediately have

$$|p_m| = \left|\frac{p^{(m)}(0)}{m!}\right| \leq |B_1|, \ m \in \mathbb{N}, \tag{2.14}$$

and

$$|q_m| = \left|\frac{q^{(m)}(0)}{m!}\right| \leq |B_1|, \ m \in \mathbb{N}. \tag{2.15}$$

Applying (2.14), (2.15) and Lemma 1.8 for the coefficients p_1, p_2, q_1 and q_2, we readily get

$$|a_2| \leq \sqrt{\frac{2|B_1|\cos\beta}{(1+\mu)(2\lambda+\mu)}}.$$

This gives the bound on $|a_2|$ as asserted in (2.1).

Next, in order to find the bound on $|a_3|$, by subtracting (2.10) from (2.8), we get

$$2(a_3 - a_2^2)(2\lambda+\mu) = e^{-i\beta}(p_2 - q_2)\cos\beta. \tag{2.16}$$

It follows from (2.13) and (2.16) that

$$a_3 = \frac{e^{-i\beta}\cos\beta(p_2+q_2)}{(1+\mu)(2\lambda+\mu)} + \frac{e^{-i\beta}(p_2-q_2)\cos\beta}{2(2\lambda+\mu)}. \tag{2.17}$$

Applying (2.14), (2.15) and Lemma 1.8 once again for the coefficients p_1, p_2, q_1 and q_2, we readily get

$$|a_3| \leq \frac{2|B_1| cos\beta}{(2\lambda+\mu)(1+\mu)}.$$

This completes the proof of Theorem 2.1. □

3. Corollaries and Consequences

In view of Remark 1.2, if we set

$$h(z) = \frac{1+Az}{1+Bz}, \quad -1 \leq B < A \leq 1, \ z \in \mathbb{U}$$

and

$$h(z) = \frac{1+(1-2\alpha)z}{1-z}, \ 0 \leq \alpha < 1, \ z \in \mathbb{U},$$

in Theorem 2.1, we can readily deduce Corollaries 3.1 and 3.2, respectively, which we merely state here without proof.

Corollary 3.1. *Let $f(z)$ given by (1.1) be in the class $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta, \frac{1+Az}{1+Bz})$, then*

$$|a_2| \leq \sqrt{\frac{2(A-B)\cos\beta}{(1+\mu)(2\lambda+\mu)}} \tag{3.1}$$

and

$$|a_3| \leq \frac{2(A-B)cos\beta}{(2\lambda+\mu)(1+\mu)}, \tag{3.2}$$

where $\beta \in (-\pi/2, \pi/2)$, $\mu \geq 0$ and $\lambda \geq 1$.

Corollary 3.2. *Let* $f(z)$ *given by (1.1) be in the class* $\mathcal{NP}_{\Sigma}^{\mu,\lambda}(\beta,\alpha)$, $0 \leq \alpha < 1$, $\mu \geq 0$ *and* $\lambda \geq 1$, *then*

$$|a_2| \leq \sqrt{\frac{4(1-\alpha)\cos\beta}{(1+\mu)(2\lambda+\mu)}} \tag{3.3}$$

and

$$|a_3| \leq \frac{4(1-\alpha)cos\beta}{(2\lambda+\mu)(1+\mu)}, \tag{3.4}$$

where $\beta \in (-\pi/2, \pi/2)$.

Remark 3.3. When $\beta = 0$ the estimates of the coefficients $|a_2|$ and $|a_3|$ of the Corollary 3.2 are improvement of the estimates obtained in [5, Theorem 3.1].

Corollary 3.4. *Let* $f(z)$ *given by (1.1) be in the class* $\mathcal{NP}_{\Sigma}^{\mu,1}(\beta,\alpha)$, $0 \leq \alpha < 1$ *and* $\mu \geq 0$, *then*

$$|a_2| \leq \sqrt{\frac{4(1-\alpha)\cos\beta}{(1+\mu)(2+\mu)}} \tag{3.5}$$

and

$$|a_3| \leq \frac{4(1-\alpha)cos\beta}{(2+\mu)(1+\mu)}, \tag{3.6}$$

where $\beta \in (-\pi/2, \pi/2)$.

Corollary 3.5. *Let* $f(z)$ *given by (1.1) be in the class* $\mathcal{NP}_{\Sigma}^{0,1}(\beta,\alpha)$, $0 \leq \alpha < 1$, *then*

$$|a_2| \leq \sqrt{2(1-\alpha)\cos\beta} \tag{3.7}$$

and

$$|a_3| \leq 2(1-\alpha)cos\beta, \tag{3.8}$$

where $\beta \in (-\pi/2, \pi/2)$.

Remark 3.6. Taking $\beta = 0$ in Corollary 3.5, the estimate (3.7) reduces to $|a_2|$ of [10, Corollary 3.3] and (3.8) is improvement of $|a_3|$ obtained in [10, Corollary 3.3].

Corollary 3.7. *Let* $f(z)$ *given by (1.1) be in the class* $\mathcal{NP}_{\Sigma}^{1,\lambda}(\beta,\alpha)$, $0 \leq \alpha < 1$ *and* $\lambda \geq 1$, *then*

$$|a_2| \leq \sqrt{\frac{2(1-\alpha)\cos\beta}{2\lambda+1}} \tag{3.9}$$

and

$$|a_3| \leq \frac{2(1-\alpha)cos\beta}{2\lambda+1}, \tag{3.10}$$

where $\beta \in (-\pi/2, \pi/2)$.

Remark 3.8. Taking $\beta = 0$ in Corollary 3.7, the inequality (3.10) improves the estimate of $|a_3|$ in [6, Theorem 3.2].

Corollary 3.9. *Let* $f(z)$ *given by (1.1) be in the class* $\mathcal{NP}_{\Sigma}^{1,1}(\beta,\alpha)$, $0 \leq \alpha < 1$, *then*

$$|a_2| \leq \sqrt{\frac{2(1-\alpha)\cos\beta}{3}} \tag{3.11}$$

and

$$|a_3| \leq \frac{2(1-\alpha)cos\beta}{3}, \tag{3.12}$$

where $\beta \in (-\pi/2, \pi/2)$.

Remark 3.10. For $\beta = 0$ the inequality (3.12) improves the estimate $|a_3|$ of [14, Theorem 2].

REFERENCES

[1] R. M. Ali, S.K.Lee, V.Ravichandran, S.Supramanian, Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions, Appl. Math. Lett. **25** (2012), no. 3, 344–351.

[2] D.A. Brannan, J.G. Clunie (Eds.), Aspects of contemporary complex analysis, Academic Press, London, 1980.

[3] D. A. Brannan, J. Clunie and W. E. Kirwan, Coefficient estimates for a class of starlike functions, Canad. J. Math. **22** (1970), 476–485.

[4] D. A. Brannan and T. S. Taha, On some classes of bi-univalent functions, Studia Univ. Babeş-Bolyai Math. **31** (1986), no. 2, 70–77.

[5] M.Çağlar, H.Orhan and N.Yağmur, Coefficient bounds for new subclasses of bi-univalent functions, Filomat, in press.

[6] B. A. Frasin and M. K. Aouf, New subclasses of bi-univalent functions, Appl. Math. Lett. **24** (2011), no. 9, 1569–1573.

[7] S.P.Goyal and P.Goswami, Estimate for initial Maclaurin coefficients of bi-univalent functions for a class defined by fractional derivatives, J. Egyptian Math. Soc., **20** (2012), 179–182.

[8] T. Hayami and S. Owa, Coefficient bounds for bi-univalent functions, *Pan Amer. Math. J.* **22** (4) (2012), 15–26.

[9] M. Lewin, On a coefficient problem for bi-univalent functions, Proc. Amer. Math. Soc. **18** (1967), 63–68.

[10] X.-F. Li and A.-P. Wang, Two new subclasses of bi-univalent functions, *Internat. Math. Forum* **7** (2012), 1495–1504.

[11] E. Netanyahu, The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$, Arch. Rational Mech. Anal. **32** (1969), 100–112.

[12] W. Rogosinski, On the coefficients of subordinate functions, Proc. London Math. Soc. (2) **48** (1943), 48–82.

[13] S. Sivaprasad Kumar, V.Kumar and V. Ravichandran, Estimates for the initial coefficients of bi-univalent functions, Preprint.

[14] H. M. Srivastava, A. K. Mishra and P. Gochhayat, Certain subclasses of analytic and bi-univalent functions, Appl. Math. Lett. **23** (2010), no. 10, 1188–1192.

[15] T.S. Taha, Topics in univalent function theory, Ph.D. Thesis, University of London, 1981.

[16] Q.-H. Xu, H.M. Srivastava, Z. Li, A certain subclass of analytic and close-to-convex functions, Appl. Math. Lett. 24 (2011) 396–401.

[17] Q.-H. Xu, Y.-C. Gui and H. M. Srivastava, Coefficient estimates for a certain subclass of analytic and bi-univalent functions, Appl. Math. Lett. **25** (2012), no. 6, 990–994.

[18] Q.-H. Xu, H.-G. Xiao and H. M. Srivastava, A certain general subclass of analytic and bi-univalent functions and associated coefficient estimate problems, Appl. Math. Comput. **218** (2012), no. 23, 11461–11465.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, ATATURK UNIVERSITY, 25240 ERZURUM, TURKEY
E-mail address: `orhanhalit607@gmail.com; horhan@atauni.edu.tr`

POST-GRADUATE AND RESEARCH DEPARTMENT OF MATHEMATICS, GOVERNMENT ARTS COLLEGE FOR MEN, KRISHNAGIRI 635001, TAMILNADU, INDIA,
E-mail address: `nmagi_2000@yahoo.co.in`

DEPARTMENT OF MATHEMATICS, L.N. GOVT COLLEGE, PONNERI, CHENNAI, TAMILNADU, INDIA
E-mail address: `balajilsp@yahoo.co.in`